Asymptotic behaviors of radial solutions for a class of semilinear elliptic equations

SHAOHUA CHEN

Department of Mathematics and Statistics, Simon Fraser University

Burnaby, B.C., Canada, V5A 1S6

GUOZHEN LU*

Department of Mathematics and Statistics, Wright State University

Dayton, Ohio 45435, USA

KEY WORDS: Semilinear equations, Pokhozhaev's identities, Oscillatory solutions, Amplitude, Periods, Asymptotic behaviors. AMS classification: 35J10, 35J70, 42B20 and 31B35.

1 Introduction

The main purpose of this paper is to investigate the asymptotic behavior of the oscillatory solutions to the semilinear equation of the form

$$\triangle u + f(|x|, u) = 0 \quad \text{in} R^n, n \geq 3. \tag{1}$$

Equation (1) has its origin from, e.g., the prescribed curvature problems in Riemannian geometry, and astrophysics (i.e., the Lane-Emden-Fowler equation and the Matukuma equation as special cases). The asymptotic behavior of the positive solutions to (1) has recently received much attentions, see e.g., [L1], [LN], [Na], etc.

However, it is well-known that the above equation (1) does not always have positive solutions or positive radial solutions. In other words, under suitable conditions, radial solutions to (1)

*The second author's research is supported in part by the National Science Foundation Grant #DMS93-15963.

must oscillate about the zero at infinite times (see e.g., [DCC], [NY], [N], etc.). Thus it becomes very interesting to know as precisely as possible the asympototic behaviors of the oscillating periods, the amplitudes of the oscilatory solutions.

In this paper, we restrict our attention to the study of radial solutions to (1). More precisely, we shall discuss the asymptotic behavior of oscilatory radial solutions for the following singular initial value problem for the semilinear elliptic equation

$$u'' + \frac{n-1}{r}u' + (br^{\mu}|u|^{q-1} + r^{\nu}|u|^{p-1})u = 0,\ u(0) = a \neq 0, u'(0) = 0, \tag{2}$$

where $b > 0, n \geq 3, \mu > -2, \nu > -2, 1 < q < \frac{n+2+2\mu}{n-2}$ and

$$p = \frac{n+2+2\nu}{n-2} = 1 + \frac{4+2\nu}{n-2} > 1. \tag{3}$$

Ni and Yotsutani [NY] have considered the equation

$$u'' + \frac{n-1}{r}u' + \sum_{i=1}^{k} c_i r^{l_i}(u^+)^{p_i} = 0,\ u(0) = u_0 > 0, u'(0) = 0, \tag{4}$$

where $p_i > 1, l_i > -2, c_i > 0, (1 \leq i \leq k)$ and $u^+ = \max[u(r), 0]$. They concluded for (4) that:

(a) if $p_i \leq (n+2+2l_i)/(n-2)$, for $1 \leq i \leq k$, with at least one inequality being a strict, then $u(r)$ has a finite zero for every u_0.

(b) if $p_i \geq (n+2+2l_i)/(n-2)$, for $1 \leq i \leq k$, then $u(r)$ is a positive solution for every u_0.

Nishihara [N] delt with the second order differential equation

$$y''(s) + f(s)|y(s)|^{p-1}y(s) = 0,$$

which is equivalent to

$$u''(r) + \frac{n-1}{r}u'(r) + k(r)|u(r)|^{p-1}u(r) = 0,$$

by the transformation $u(r) = y(s)$ and $s = (n-2)^{n-2}r^{-(n-2)}$. Under suitable assumptions on f(s), Nishihara proved that the solution $y(s)$ is nonoscillatory.

W. R. Derrick, S. Chen, and J. A. Cima [DCC] recently discussed the following equation when the restoring function $f(|x|, u)$ is independent of the radial variable $|x|$:

$$u'' + \frac{n-1}{r}u' + (b|u|^{q-1} + |u|^{p-1})u = 0, u(0) \neq 0, u'(0) = 0, \tag{5}$$

$1 < q < p = \frac{n+2}{n-2}$. They proved that

$$\lim_{j\to\infty} t_j^{\frac{2n-2}{q+3}}|u(t_j)| = c \ \ and \ \ \frac{r_{j+1} - r_j}{r_{j+1}^{\frac{(n-1)(q-1)}{q+3}}} \geq c,$$

where c is a positive constant, t_j and r_j are the j^{th} local extremum and zeros of the solution $u(r)$, respectively. This shows that the period between two consecutive zeros $r_{j+1} - r_j$ has a lower bound of the order $r_{j+1}^{\frac{(n-1)(q-1)}{q+3}}$ as r_j goes to ∞.

This paper generalizes and improves the results of [DCC] in several ways. First of all, we allow the restoring fnction $f(|x|, u)$ to depend on $|x|$ rather than to be a function of u only. Secondly, we provide an exact asymptotic behavior of the period between any two consecutive zeros r_j and r_{j+1}. Namely, we also show an upper bound of the period, which is of the same order as the lower bound. The precise statement is given in Theorem 2 below. We also derive in this paper the precise asymptotic estimate for the energy associated to the equation (2) defined by (8) which is of independent interest, see Theorem 1 for details.

2 Preliminary

We will often use the following Pokhozhaev's second identity (see [Po]):

$$\begin{aligned} r^{k+1}[\frac{1}{2}u'^2(r) + F(r, u(r))] + \alpha r^k u'(r)u(r) + \frac{\alpha}{2}(n-1-k)r^{k-1}u^2(r) \\ +(n-1-\frac{k+1}{2}-\alpha)\int_{r_0}^r u'^2(s)s^k ds - \alpha\frac{(n-1-k)(k-1)}{2}\int_{r_0}^r u^2(s)s^{k-2}ds \\ +\int_{r_0}^r [\alpha u(s)f(s,u(s)) - (k+1)F(s,u(s)) - sF_s(s,u(s))]s^k ds = c_0 \end{aligned} \tag{6}$$

for any k and $r_0 > 0$, where

$$c_0 = r_0^{k+1}[\frac{1}{2}u'^2(r_0) + F(r_0, u(r_0))] + \alpha r_0^k u'(r_0)u(r_0) + \frac{\alpha}{2}(n-1-k)r_0^{k-1}u^2(r_0)$$

and

$$F(r, u) = \int_0^u f(r, v)dv,$$

and $u(r)$ is a solution of

$$u''(r) + \frac{n-1}{r}u'(r) + f(r, u) = 0. \tag{7}$$

Let $u(r)$ be a solution of (2) and define the energy function

$$Q(r, u(r)) = \frac{1}{2}u'^2(r) + F(r, u(r)) = \frac{1}{2}u'^2(r) + \frac{br^\mu|u(r)|^{q+1}}{q+1} + \frac{r^\nu|u(r)|^{p+1}}{p+1}. \tag{8}$$

Suppose that the solution $u(r)$ of (2) oscillates about the zero a finite number of times and has a local maximum at r_1 for which $u(r) > 0$ for all $r > r_1$. We call such solutions eventually positive solutions.

Lemma 1 *The problem (2) has no eventually positive or negative solutions.*

Remark 1: W. Ni and S. Yotsutani [NY] proved that problem (2) has a finite number of zeros for every $a > 0$. The techniques of [NY] can be adapted to prove our Lemma 1. However, we provide here an alternate proof for the sake of completeness.

Proof of Lemma 1: Since the case for eventually negative solutions follows trivially by setting $v = -u$, we only prove that there is no eventually positive solutions. Suppose that the conclusion were not true, then there would exist some $r_1 \geq 0$ such that $u'(r_1) = 0$ and $u(r) > 0$ for all $r \geq r_1$.

Note

$$u'(r) = -\frac{1}{r^{n-1}} \int_{r_1}^{r} [bs^{\mu} u^q(s) + s^{\nu} u^p(s)] s^{n-1} ds < 0 \quad for\ r > r_1, \tag{9}$$

which implies that $u(r)$ is decreasing for $r > r_1$. Then

$$u'(r) \leq -\frac{b}{r^{n-1}} u^q(r) \int_{r_1}^{r} s^{n-1+\mu} ds = -\frac{b}{r^{n-1}} \cdot \frac{u^q(r)}{n+\mu} (r^{n+\mu} - r_1^{n+\mu}),$$

or

$$\frac{du(r)}{u^q(r)} \leq -\frac{b}{n+\mu} (r^{\mu+1} - r_1^{n+\mu} r^{1-n}) dr.$$

Integrating from r_1 to r we get

$$\frac{1}{q-1} \left(\frac{1}{u^{q-1}(r_1)} - \frac{1}{u^{q-1}(r)} \right) \leq -\frac{b}{n+\mu} \left(\frac{1}{\mu+2} r^{\mu+2} + \frac{1}{n-2} r_1^{n+\mu} r^{2-n} - \frac{n+\mu}{(n-2)(\mu+2)} r_1^{\mu+2} \right),$$

or

$$\frac{1}{u^{q-1}(r_1)} + \frac{b(q-1)}{n+\mu} \left(\frac{1}{\mu+2} r^{\mu+2} + \frac{1}{n-2} r_1^{n+\mu} r^{2-n} - \frac{n+\mu}{(n-2)(\mu+2)} r_1^{\mu+2} \right) \leq \frac{1}{u^{q+1}(r)}.$$

Thus

$$u(r) \leq c r^{-\frac{\mu+2}{q-1}} \tag{10}$$

for $r > r_2$ with sufficiently large $r_2 \geq r_1$.

Inserting (10) into (9) we obtain

$$\begin{aligned} |u'(r)| \quad &\leq \frac{c}{r^{n-1}} + \frac{c}{r^{n-1}} \int_{r_1}^{r} s^{n-1+\mu-\frac{(\mu+2)q}{q-1}} ds + \int_{r_1}^{r} s^{n-1+\nu-\frac{(\mu+2)p}{q-1}} ds \\ &\leq c \cdot \max \left(r^{-\frac{\mu+2}{q-1}-1}, r^{\nu+1-\frac{(\mu+2)p}{q-1}}, r^{-(n-1)} \right). \end{aligned} \tag{11}$$

Applying (6) with $k = n-1$, $r_0 = 0$ and $\alpha = \frac{n-2}{2}$ we get

$$r^n \left(\frac{1}{2} u'^2(r) + F(r, u(r)) \right) + \frac{n-2}{2} r^{n-1} u'(r) u(r) = \frac{n+2+2\mu-(n-2)q}{2(q+1)} \int_0^r |u(s)|^{q+1} s^{n-1+\mu} ds \tag{12}$$

Using (10) and (11), we derive

$$r^n\left(\frac{1}{2}u'^2(r)+\frac{br^\mu|u(r)|^{q+1}}{q+1}\right)\le c\cdot\max\left(r^{-\frac{n+2+2\mu-(n-2)q}{q-1}},r^{n+2(\nu+1-\frac{(\mu+2)p}{q-1})},r^{-(n-2)}\right)$$

and

$$r^{n+\nu}\frac{|u(r)|^{p+1}}{p+1}\le c\,r^{-\frac{n+2+2\mu-(n-2)q}{(q-1)(n-2)}\cdot(n+\nu)}.$$

Letting $r\to\infty$ in (12) we get a contradiction, since the left side of (12) is nonpositive, while the right side is a positive number. Thus Lemma (1) follows.

3 Main theorems and their proofs

By Lemma (1), the solutions of (2) must oscillate about zero infinitely many times. Let r_j be the zeros of $u(r)$ and t_j be the local extremum points of $u(r)$ with $r_j<t_j<r_{j+1}$. Now we will study the amplitudes and the periods of the oscillations.

Theorem 1 *There exists a positive constant c_1 such that*

$$\lim_{r\to\infty}r^{\frac{2(n-1)(q+1)-2\mu}{q+3}}Q(r,u(r))=c_1. \tag{13}$$

Consequently,

$$\lim_{j\to\infty}t_j^{\frac{2n-2+\mu}{q+3}}|u(t_j)|=\left[\frac{(q+1)c_1}{b}\right]^{\frac{1}{q+1}}\ \textit{and}\ \lim_{j\to\infty}r_j^{\frac{(n-1)(q+1)-\mu}{q+3}}|u'(r_j)|=\sqrt{2c_1}. \tag{14}$$

Proof: Applying (6) with $\alpha=\frac{2n-2+\mu}{q+3}$ and

$$k=\frac{2(n-1)(q+1)-2\mu}{q+3}-1=n-1-\frac{n+2+2\mu-(n-2)q}{q+3}<n-1, \tag{15}$$

we have

$$\begin{aligned}&r^{k+1}\left(\frac{1}{2}u'^2(r)+F(r,u(r))\right)+\alpha r^k u'(r)u(r)+c_2r^{k-1}u^2(r)\\+c_3&\int_{r_0}^r|u(s)|^{p+1}s^{k+\nu}ds+c_3'\int_{r_0}^r s^\mu|u|^{q+1}ds=c_4\int_{r_0}^r u^2(s)s^{k-2}ds+c_0,\end{aligned} \tag{16}$$

or

$$\begin{aligned}&\frac{1}{4}r^{k+1}u'^2(r)+r^{k+1}\left(\frac{b}{q+1}r^\mu|u(r)|^{q+1}+\frac{1}{p+1}r^\nu|u(r)|^{p+1}\right)\\&+\frac{1}{4}\left[ru'(r)+2\alpha u(r)\right]^2+c_3\int_{r_0}^r|u(s)|^{p+1}s^{k+\nu}ds+c_3'\int_{r_0}^r s^{k+\mu}|u|^{q+1}ds\\&=c_4\int_{r_0}^r u^2(s)s^{k-2}ds+c_0+(\alpha^2-c_2)r^{k-1}u^2(r),\end{aligned} \tag{17}$$

where c_0 is defined as in (6),

$$c_3' = b\left(\alpha - \frac{k+1}{q+1} - \frac{\mu}{q+1}\right) = 0,$$

$$c_2 = \frac{(2n-2+\mu)(n+2+2\mu-(n-2)q)}{2(q+3)^2},\ c_4 = c_2(k-1),$$

and

$$c_3 = \alpha - \frac{k+1}{p+1} - \frac{\nu}{p+1} = \frac{2(n-1)(p-q)+\mu(p+3)-\nu(q+3)}{(q+3)(p+1)}.$$

Since

$$\begin{aligned}
&2(n-1)(p-q)+\mu(p+3)-\nu(q+3)-2[n+2+2\mu-(n-2)q] \\
&= [2(n-1)+\mu]p - 2(n-1)q + 3\mu - \nu q - 3\nu - 2(n+2) - 4\mu + 2q(n-2) \\
&= [2(n-1)+\mu]\frac{n+2+2\nu}{n-2} - \mu - 2q - \nu q - 3\nu - 2(n+2) \\
&= (2+\nu)\frac{n+2+2\mu-(n-2)q}{n-2} > 0, \qquad (18)
\end{aligned}$$

we then get $c_3 > 0$. By Holder's inequality with two congugate exponents $\frac{p+1}{2}$ and $\frac{p+1}{p-1}$ we have

$$|c_4|u^2(s)s^{k-2} = (u^2(s)s^{\frac{2(k+\nu)}{p+1}})(|c_4|s^{k-2-\frac{2(k+\nu)}{p+1}}) \le c_3|u(s)|^{p+1}s^{k+\nu} + c\,s^{\beta}, \qquad (19)$$

where

$$\beta = \left(k-2-\frac{2k+2\nu}{p+1}\right)\cdot\frac{p+1}{p-1} = k-2-\frac{4+2\nu}{p-1} = k-n+\frac{(n-2)p-n-2-2\nu}{p-1} = k-n < -1$$

by (15). Using (19), we also get

$$|\alpha^2 - c_2|r^{k-1}u^2(r) = |\alpha^2-c_2|r\cdot[u^2(r)r^{k-2}] \le \frac{1}{2(p+1)}|u(r)|^{p+1}r^{k+1+\nu} + c\,r^{\beta+1}. \qquad (20)$$

Substituting (19) and (20) into (17) yields

$$\begin{aligned}
&\frac{1}{4}r^{k+1}u'^2(r) + r^{k+1}\left[\frac{b}{q+1}r^{\mu}|u(r)|^{q+1} + \frac{1}{p+1}r^{\nu}|u(r)|^{p+1}\right] + c_3\int_{r_0}^{r}|u(s)|^{p+1}s^{k+\nu}ds \\
&\le c_3\int_{r_0}^{r}|u(s)|^{p+1}s^{k+\nu}ds + c\int_{r_0}^{r}s^{\beta}ds + \frac{1}{2(p+1)}r^{k+1+\nu}|u(r)|^{p+1} + cr^{\beta+1} + c_0. \qquad (21)
\end{aligned}$$

Thus we have simultaneously proved that

$$r^{k+1}Q(r,u(r)) \le c,\ |u(r)| \le c\,r^{-\frac{2n-2+\mu}{q+3}}\ \textit{and}\ |u'(r)| \le c\,r^{-\frac{(n-1)(q+1)-\mu}{q+3}}. \qquad (22)$$

Consequently we have from (22)

$$|\alpha r^k u(r)u'(r)| \le c\,r^{\frac{2(n-1)(q+1)-2\mu}{q+3}-1}\cdot r^{-\frac{2n-2+\mu}{q+3}}\cdot r^{-\frac{(n-1)(q+1)-\mu}{q+3}} = c\,r^{-\frac{n+2+2\mu-(n-2)q}{q+3}}, \qquad (23)$$

$$r^{k-1}u^2(r) \le c\, r^{\frac{2(n-1)(q+1)-2\mu}{q+3}-2} \cdot r^{-\frac{2(2n-2+\mu)}{q+3}} = c\, r^{-\frac{2[n+2+2\mu-(n-2)q]}{q+3}}, \tag{24}$$

Letting $r \to \infty$ in (16) and using (23) and (24) we derive

$$\lim_{r\to\infty} r^{k+1}Q(r,u(r)) = c_4 \int_{r_0}^{\infty} u^2(s)s^{k-2}ds - c_3 \int_{r_0}^{\infty} |u(s)|^{p+1}s^{k+\nu}ds + c_0. \tag{25}$$

Now we claim that

$$\lim_{r\to\infty} r^{k+1}Q(r,u(r)) = c_1 > 0. \tag{26}$$

Suppose otherwise, then $c_1 = 0$, since $Q(r,u(r)) > 0$ for all $r > 0$. Set

$$\lambda = \frac{n+2+2\mu-(n-2)q}{q+3} - \epsilon_0 > 2(q+1)\epsilon_0 > 0$$

for small ϵ_0. By L'Hospital's rule, (22)-(24) and (18) we will get

$$\begin{aligned}
&\left|\lim_{r\to\infty} \frac{c_4 \int_{r_0}^{r} u^2(s)s^{k-2}ds - c_3 \int_{r_0}^{r} |u(s)|^{p+1}s^{k+\nu}ds + c_0}{r^{-\lambda}}\right| \\
&\le \frac{c}{\lambda}\left(\lim_{r\to\infty} u^2(r)r^{k-1+\lambda} + \lim_{r\to\infty} |u(r)|^{p+1}r^{k+1+\lambda+\nu}\right) \\
&\le \frac{c}{\lambda}\left(\lim_{r\to\infty} r^{\lambda-\frac{2(n+2+2\mu-(n-2)q)}{q+3}} + \lim_{r\to\infty} r^{\lambda-\frac{2(n-1)(p-q)+(p+3)\mu-\nu(q+3)}{q+3}}\right) \\
&= 0.
\end{aligned} \tag{27}$$

Thus, multiplying (16) by r^{λ} and using (23), (24) and (27) we get

$$\lim_{r\to\infty} r^{k+1+\lambda}Q(r,u(r)) = 0 \textit{ or } |u(r)| \le c\, r^{-\frac{k+1+\lambda+\mu}{q+1}} \le c\, r^{-\frac{2n-2+\mu}{q+3}-\frac{\lambda}{q+1}}. \tag{28}$$

Consequently, we have by (28), (22) and (18),

$$|\alpha r^k u(r)u'(r)| \le c\, r^{-\frac{n+2+2\mu-(n-2)q}{q+3}-\frac{\lambda}{q+1}} \le c\, r^{-\frac{2(n+2+2\mu-(n-2)q)}{q+3}-2\epsilon_0} \tag{29}$$

and

$$|u(r)|^{p+1}r^{k+1+\nu} \le c\, r^{-\frac{2(n-1)(p-q)+(p+3)\mu-\nu(q+3)}{q+3}-\frac{\lambda(p+1)}{q+1}} \le cr^{-\frac{n+2+2\mu-(n-2)q}{q+3}-2\epsilon_0}. \tag{30}$$

We can repeat the process in (27) by using

$$\lambda_1 = \frac{n+2+2\mu-(n-2)q}{q+3} + \epsilon_0$$

to obtain

$$\lim_{r\to\infty} r^{k+1+\lambda_1}Q(r,u(r)) = 0. \tag{31}$$

On the other hand, using (12) with $r = r_j$ and $u(r_j) = 0$, we get

$$r_j^n Q(r_j,u(r_j)) = \frac{n+2+2\mu-(n-2)q}{2(q+1)} \int_0^{r_j} |u(s)|^{q+1}s^{n-1+\mu}ds > 0$$

which contradicts (31) since $k+1+\lambda_1 = n+\epsilon_0$. Hence $c_1 > 0$ in (26) and the proof is complete.

Theorem 2 *There exist two positive constants $0 < c_2 < c_3$ such that*

$$c_2 \leq \frac{r_{j+1} - r_j}{r_{j+1}^{\frac{(n-1)(q-1)-2\mu}{q+3}}} \leq c_3. \tag{32}$$

Proof: Since $u'(t_j) = 0$ and $u(r)$ does not change sign in $[t_j, r_{j+1}]$, we have by (14), (9) and (10) for j large enough,

$$\begin{aligned} \sqrt{c_1} \quad & \leq r_{j+1}^{\frac{(n-1)(q+1)-\mu}{q+3}} |u'(r_{j+1})| \\ & = r_{j+1}^{-\frac{2(n-1)+\mu}{q+3}} \int_{t_j}^{r_{j+1}} (bs^\mu |u(s)|^q + s^\nu |u(s)|^p) \, s^{n-1} ds \\ & \leq c \, r_{j+1}^{-\frac{2(n-1)+\mu}{q+3}} \int_{t_j}^{r_{j+1}} s^{n-1+\mu-\frac{(2n-2+\mu)q}{q+3}} ds \\ & \leq c \, r_{j+1}^{-\frac{2(n-1)+\mu}{q+3}} \int_{t_j}^{r_{j+1}} s^{\frac{(n-1)(3-q)+3\mu}{q+3}} ds. \end{aligned} \tag{33}$$

If $(n-1)(3-q) + 3\mu \geq 0$, then the integrand is monotone increasing so that

$$\sqrt{c_1} \leq c \, r_{j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} \int_{t_j}^{r_{j+1}} ds \leq c \, r_{j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} (r_{j+1} - r_j).$$

If $(n-1)(3-q) + 3\mu < 0$, since

$$\frac{(n-1)(3-q) + 3\mu}{q+3} + 1 = \frac{3n - (n-2)q + 3\mu}{q+3} > 0$$

we can calculate (33) directly and get

$$\begin{aligned} \sqrt{c_1} \quad & \leq \frac{c(q+3)}{3n - q(n-2) + 3\mu} r_{j+1}^{-\frac{2(n-1)+\mu}{q+3}} \left(r_{j+1}^{\frac{(n-1)(3-q)+3\mu}{q+3}+1} - t_j^{\frac{(n-1)(3-q)+3\mu}{q+3}+1} \right) \\ & = c \, r_{j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} \left(r_{j+1} - r_{j+1}^{-\frac{(n-1)(3-q)+3\mu}{q+3}} \cdot t_j^{\frac{(n-1)(3-q)+3\mu}{q+3}+1} \right) \\ & \leq c \, r_{j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} (r_{j+1} - t_j) \\ & \leq c \, r_{j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} (r_{j+1} - r_j). \end{aligned} \tag{34}$$

Thus we have proved the first inequality in (32).

Now let us prove the second inequality in (32). First suppose $u(t_{2j}) > 0$ for $j = 1, 2, \cdots$. Let $s_{2j} \in (t_{2j}, r_{2j+1})$ such that

$$u'^2(s_{2j}) = 2F(s_{2j}, u(s_{2j})) \text{ and } u'^2(r) > 2F(r, u(r)), \tag{35}$$

for $r \in (s_{2j}, r_{2j+1})$. Note that such s_{2j} must exist because if we set

$$g(r) = u'^2(r) - 2F(r, u(r)) = u'^2(r) - 2\left(br^\mu \frac{|u(r)|^{q+1}}{q+1} + r^\nu \frac{|u(r)|^{p+1}}{p+1} \right),$$

then $g(t_{2j}) = -2F(t_{2j}, u(t_{2j})) < 0$ and $g(r_{2j+1}) = u'^2(r_{2j+1}) > 0$. Thus

$$s_{2j} = \sup_{r \in (t_{2j}, r_{2j+1})} \{r \,|u'^2(r) = 2F(r, u(r))\}.$$

Using (13) and (35) we have

$$s_{2j}^{k+1} Q(s_{2j}, u(s_{2j})) = 2s_{2j}^{k+1} F(s_{2j}, u(s_{2j})) \to c_1 \; as \; j \to \infty,$$

which implies that

$$s_{2j}^{\frac{2(n-1)+\mu}{q+3}} u(s_{2j}) \to \left(\frac{(q+1)c_1}{2b}\right)^{\frac{1}{q+1}} = c_0 \; as \; j \to \infty. \tag{36}$$

Using (13) and (35) again, we get

$$\frac{c_1}{2} < r^{k+1} Q(r, u(r)) \leq r^{k+1} u'^2(r),$$

for $r \in (s_{2j}, r_{2j+1})$ and j large enough, which implies that (notice $u'(r) < 0$ in (t_{2j}, r_{2j+1}))

$$r^{\frac{(n-1)(q+1)-\mu}{q+3}}(-u'(r)) = r^{\frac{k+1}{2}}(-u'(r)) \geq \sqrt{\frac{c_1}{2}}. \tag{37}$$

Multiplying both sides of (37) by $r^{-\frac{(n-1)(q-1)-2\mu}{q+3}}$ and adding a like term to both sides, gives

$$\begin{aligned} & -\frac{2(n-1)+\mu}{q+3} u(r) r^{\frac{2(n-1)+\mu}{q+3}-1} - u'(r) r^{\frac{2(n-1)+\mu}{q+3}} \\ \geq \quad & \sqrt{\frac{c_1}{2}} r^{-\frac{(n-1)(q-1)-2\mu}{q+3}} - \frac{2(n-1)+\mu}{q+3} u(r) r^{\frac{2(n-1)+\mu}{q+3}-1}, \end{aligned}$$

or

$$-\frac{d}{dr}\left(u(r) r^{\frac{2(n-1)+\mu}{q+3}}\right) \geq r^{-\frac{(n-1)(q-1)-2\mu}{q+3}} \left(\sqrt{\frac{c_1}{2}} - \frac{2(n-1)+\mu}{q+3} u(r) r^{\frac{2(n-1)+\mu}{q+3}} \cdot r^{-\frac{n+2-(n-2)q+2\mu}{q+3}}\right). \tag{38}$$

Since $u(r) r^{\frac{2(n-1)+\mu}{q+3}}$ is bounded by (22) and $r^{-\frac{n+2+2\mu-(n-2)q}{q+3}} \to 0$ as $r \to \infty$, we can assume that

$$\left(\sqrt{\frac{c_1}{2}} - \frac{2(n-1)+\mu}{q+3} u(r) r^{\frac{2(n-1)+\mu}{q+3}} \cdot r^{-\frac{n+2+2\mu-(n-2)q}{q+3}}\right) \geq \frac{\sqrt{c_1}}{2},$$

for $r \in (s_{2j}, r_{2j+1})$ and j sufficiently large. Thus (38) becomes

$$-\frac{d}{dr}\left(u(r) r^{\frac{2(n-1)+\mu}{q+3}}\right) \geq \frac{\sqrt{c_1}}{2} r^{-\frac{(n-1)(q-1)-2\mu}{q+3}} \tag{39}$$

Integrating (39) from s_{2j} to r_{2j+1} yields, by (36),

$$2c_0 \geq s_{2j}^{\frac{2(n-1)+\mu}{q+3}} u(s_{2j}) \geq \begin{cases} \frac{\sqrt{c_1}}{2} r_{2j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} (r_{2j+1} - s_{2j}), & \text{if } (n-1)(q-1) \geq 2\mu; \\ \frac{\sqrt{c_1}}{2} s_{2j}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} (r_{2j+1} - s_{2j}), & \text{if } (n-1)(q-1) < 2\mu. \end{cases} \tag{40}$$

Noticing that $u(r)$ is a decreasing function on (t_{2j}, r_{2j+1}), it follows that by (35) and (9),

$$\begin{aligned} & 2\sqrt{b}s_{2j}^{\frac{\mu}{2}}u^{\frac{q+1}{2}}(s_{2j}) \geq \sqrt{2F(s_{2j}, u(s_{2j}))} = -u'(s_{2j}) \\ \geq \quad & s_{2j}^{-(n-1)}\int_{t_{2j}}^{s_{2j}} bs^{\mu}u^q(s)s^{n-1}ds \geq bs_{2j}^{-(n-1)}u^q(s_{2j})\int_{t_{2j}}^{s_{2j}} s^{n-1+\mu}ds \\ = \quad & bs_{2j}^{\mu}u^q(s_{2j})\int_{t_{2j}}^{s_{2j}} ds = bs_{2j}^{\mu}u^q(s_{2j})(s_{2j}-t_{2j}), \end{aligned}$$

or by (36)

$$\begin{aligned} 2 \geq \sqrt{b}s_{2j}^{\frac{\mu}{2}}u^{\frac{q-1}{2}}(s_{2j})(s_{2j}-t_{2j}) &\geq \frac{\sqrt{b}}{2}c_0^{\frac{q-1}{2}}s_{2j}^{-\frac{2(n-1)+\mu}{q+3}\cdot\frac{q-1}{2}+\frac{\mu}{2}}(s_{2j}-t_{2j}) \\ &\begin{cases} = \frac{\sqrt{b}}{2}c_0^{\frac{q-1}{2}}s_{2j}^{-\frac{(n-1)(q-1)-2\mu}{q+3}}(s_{2j}-t_{2j}) & \text{if } (n-1)(q-1) < 2\mu; \\ \geq \frac{\sqrt{b}}{2}c_0^{\frac{q-1}{2}}r_{2j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}}(s_{2j}-t_{2j}) & \text{if } (n-1)(q-1) \geq 2\mu. \end{cases} \end{aligned} \tag{41}$$

If $(n-1)(q-1) < 2\mu$ then by (40) and (41), $r_{2j+1} - t_{2j} \to 0$ as $j \to \infty$, which implies

$$s_{2j}^{-\frac{(n-1)(q-1)-2\mu}{q+3}} \geq \frac{1}{2}r_{2j+1}^{-\frac{(n-1)(q-1)-2\mu}{q+3}}, \tag{42}$$

for j sufficiently large.

Combining (40) - (42) we thus have proved the second inequality of (32) in the inteval (t_{2j}, r_{2j+1}).

We still need to prove the similar inequality in the interval $[r_{2j}, t_{2j}]$. Notice that

$$u'(t_{2j}) = -\frac{1}{t_{2j}^{n-1}}\int_{t_{2j-1}}^{t_{2j}} f(s, u(s))s^{n-1}ds = 0,$$

we have

$$u'(r) = -\frac{1}{r^{n-1}}\int_{t_{2j-1}}^{r} f(s, u(s))s^{n-1}ds - \frac{1}{r^{n-1}}\int_{r}^{t_{2j}} f(s, u(s))s^{n-1}ds,$$

for $r \in (r_{2j}, t_{2j})$. Then we can obtain the second inequality of (32) in the inteval (r_{2j}, t_{2j}) by similar arguments. Hence we have proved the second inequality of (32) when $u(t_{2j}) > 0$. Similarly we can prove the result when $u(t_{2j+1}) < 0$.

Remark 2: From the conclusion of Theorem 2, we notice the very interesting phenomena, that is, if $(n-1)(q-1) - 2\mu > 0$ then oscillatory period of the solution becomes longer and longer while if $(n-1)(q-1) - 2\mu < 0$ it becomes shorter and shorter. Especially, when $(n-1)(q-1) - 2\mu = 0$, it changes between two constants. For $n = 3$, $\mu = q-1$, the behaviors of solutions of (2) are very similar to that of the following linear equation

$$u'' + \frac{2}{r}u' + Au = 0, \ u(0) = a \neq 0, u'(0) = 0.$$

The solution is $u = \frac{a}{\sqrt{A}r}\sin\sqrt{A}r$ because the order of the amplitude of two solutions is the same (all are $\frac{1}{r}$).

References

[DCC] W. R. Derrick, S. Chen, and J. A. Cima, *Oscillatory radial solutions of semilinear elliptic equations*, submitted.

[L1] Yi Li, *Asymptotic behavior of positive solutions of equation* $\triangle u + K(x)u^p = 0$ *in* R^n, J. Differential Equations, 95 (1992), 304-330.

[LN] Yi Li and W.M. Ni, *On the asymptotic behavior and radial symmetry of positive solutions of semilinear elliptic equations in* R^n*. I. Asympototic Behavior,* Arch. Rational Mech. Anal., 118 (1992), 195-222.

[Na] M. Naito, *Asymptotic behavior of solutions of second order differential equations with integral coefficients*, Trans. Amer. Math. Soc., 282 (1984), 577-588.

[N] K. Nishihara, *Asymptotic behaviors of solutions of second order differential equations*, J. Math. Anal. Appl., 189(1995), 424-441.

[NY] W. -N. Ni and S. Yotsutani, *Semilinear elliptic equations of Matukuma-type and releted topics*, Japan J. Applied Math., 5(1988), 1-32.

[Po] S. I. Pokhozhaev, *On the asymptotics of entire radial solutions of quasilinear elliptic equations*, Soviet Math. Dokl., 44(1992), 548-553.